PRINCE WILLIAM COUNTY TURF CARE CALENDAR FOR WARM-SEASON GRASSES

Bermudagrass, Zoysia

Virginia Cooperative Extension - Prince William County Unit 8033 Ashton Ave, Suite 105 Manassas, VA 20109-8202

	J	an	Fe	eb	M	lar	A	pr	М	ay	Ju	ın	J	ul	Αι	ug	S	ер	С	ct	N	οv	De	ec
Soil Test (every 2 - 3 years)																								
Adjust pH (lime/sulfur)																								
Top dress with fine textured compost																								
Core aerate																								
Fertilize*																								
Mow																								
Over-seeding (if needed)**																								
Sodding damaged areas (if needed)																								
Weed Control:	\top																					П		
Summer Grasses - Pre-emergent (if needed)																								
Violet, Ground Ivy, Oxalis																								
Lespedeza, Spurge																								
Chickweed, Dandelion, Clover, Plantain***																								
Onion/Garlic Bittercress***																								
Insect Control:																								
Chinch bugs (if needed)																							П	П
Sod webworms (if needed)																								
White grubs (if needed)																								
Disease Control: ****	\top		П																			П	П	

For **Bermudagrass**, nitrogen can be applied 3-4 times at 30 day intervals through the middle of August. Most home lawns do not need more than 2 lb. N per 1,000 sf per year.

For **Zoysia grass**, fertilize once in late May/early June and again in early-mid July. Zoysia should not receive more than 2 lb. N per 1,000 sf per year.

For specific questions, call the Extension Horticulture Help Desk, 703-792-7747 or email us at mastergardener@pwcgov.org.

Virginia Cooperative Extension is a partnership of Virginia Tech, Virginia State University, the U.S. Department of Agriculture, and local governments. Its programs and employment are open to all, regardless of age, color, disability, gender, gender identity, gender expression, national origin, political affiliation, race, religion, sexual orientation, genetic information, military status, or any other basis protected by law.

^{*} Apply at a rate of no more than 1.0 lb. N per 1,000 sf per application for fertilizer with at least 15% slow release nitrogen or 0.9 lb, N per 1,000 sf per application for fertilizer with less than 15% slow release nitrogen.

^{**} Warm season grass seed is difficult to obtain and repairs are made with plugs or sod.

^{***} The timing of herbicides for these broad leaf weeds reflects when these plants most active in moving chemicals to their roots and when post emergents are effective anytime these weeds are actively growing. Please contact us for weeds not listed above.

^{****} Most common turf diseases are largely comestic and fungicides are not recommended for home lawns in most cases.